

The field of Lodén 112

N. Kaltcheva¹ and V. Golev²

¹ Department of Physics and Astronomy, University of Wisconsin Oshkosh, 800 Algoma Blvd., Oshkosh, WI 54901, USA (kaltchev@uwosh.edu)

² Department of Astronomy, Faculty of Physics, St Kliment Ohridski University of Sofia, 5 James Bourchier Blvd., BG-1164 Sofia, Bulgaria (valgol@phys.uni-sofia.bg)

Abstract

Based on the available *uvby* β photometry of OB stars in the longitude range 281° to 285° in the Galactic disk, we identify a feature of young stars at 1630 ± 82 pc, that is probably connected to the compact cluster candidate Lodén 112 and the open cluster IC 2581. This feature seems to be spatially correlated to RCW 48 and RCW 49 and several other smaller HII regions.

1 Introduction

The field between the OB groups in Vela ($262^\circ < l < 268^\circ$) and Car OB1 ($284^\circ < l < 288^\circ$) is not known to be dominated by any prominent OB association (see [2], [5]). The longitude range 283° - 284° is thought to correspond to a tangential direction of a large segment of the Carina arm in both the 3- and 4-arm models of the grand design of the Milky Way [6]. Our study is based on *uvby* β photometric distances and provides new insights on the apparent groupings and layers toward $281^\circ < l < 285^\circ$ in the Galactic disk. This field clearly stands apart from the extended HII features toward Car OB1 and contains several prominent HII regions.

2 Results

Our sample includes a number of OB stars with very similar luminosities according to the $[c_1]$ vs. $[m_1]$ classification diagram [7] (Fig. 1, left). Among them 5 stars belong to the poor but very compact cluster candidate Lodén 112. Two other stars are assigned to the relatively young cluster IC 2581, and the rest are apparent field stars. According to the *uvby* β photometric distances all OB stars in this sample show a very small spread in distance,




Figure 1: Left: the classification photometric diagram for the OB stars in the sample. Right: same stars overplotted on the distribution of the HII emission in the region [1] obtained via *SkyView* VO interface [4].

thus forming a structure at 1630 ± 80 pc. This new estimate is significantly smaller than the presently adopted distance of 2500 pc to Lodén 112 (WEBDA database). These OB-stars are located toward the prominent HII regions RCW 48 and RCW 49 and several smaller HII fields and could represent a new OB association at coordinates $282^\circ < l < 285^\circ$, $-2^\circ < b < 2^\circ$ (Fig. 1, right), containing the compact cluster candidate Lodén 112 and IC 2581 (more details can be found in [3]).

Acknowledgments

This work is supported by the National Science Foundation grant AST-0708950. N.K. acknowledges support from the SNC Endowed Professorship at the University of Wisconsin Oshkosh. V.G. acknowledges support by the Bulgarian National Science Research Fund grants DO 02-85/2008 and DO 02-362/2008. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. We acknowledge the use of NASA’s *SkyView* facility (<http://skyview.gsfc.nasa.gov>).

References

- [1] Finkbeiner, D. P. 2003, *ApJS*, 146, 407
- [2] Humphreys, R. M. 1978, *AJSS*, 38, 309
- [3] Kaltcheva, N., & Golev, V. 2011, in preparation
- [4] McGlynn, T., Scollick, K., White, N. 1998, in *New Horizons from Multi-Wavelength Sky Surveys*, Proc. of the 179th Symp. of the IAU, held in Baltimore, USA Aug 26-30, 1996, Kluwer Academic Publishers, edited by B. J. McLean, D. A. Golombek, J. J. E. Hayes, & H. E. Payne, p. 465
- [5] Melnik, A. M., & Efremov, Yu. N. 1995, *AstL*, 21, 10
- [6] Russeil, D. 2003, *A&A*, 397, 133
- [7] Strömgren, B. 1966, *ARA&A*, 4, 433